


# DOXY-PEP

DOXYCYCLINE POST-EXPOSURE PROPHYLAXIS

LA LGBT CENTER + SOUTHERN CALIFORNIA HIV/AIDS POLICY RESEARCH CENTER

EFFECTIVENESS  
IN REDUCING STIs

AMONG 1,115 PATIENTS WITH 6 MONTHS OR MORE OF DOXY-PEP USE




\* NOT DRAWN TO SCALE

24 - 72 HOURS  
DOXY-PEP

## LOS ANGELES COUNTY DEPARTMENT OF PUBLIC HEALTH RECOMMENDS

Taking a 200-mg dose of doxycycline within 24 to 72 hours of condomless sex to reduce the chance of getting a bacterial STI, such as **chlamydia, gonorrhea, syphilis**


### DOXY-PEP IS RECOMMENDED FOR:



GAY, BISEXUAL, AND OTHER MEN  
WHO HAVE SEX WITH MEN  
WHO HAVE HAD A HISTORY OF 1  
BACTERIAL STI IN THE PAST 12 MONTHS



TRANSGENDER WOMEN  
WHO HAVE HAD A HISTORY OF 1  
BACTERIAL STI IN THE PAST 12 MONTHS



INDIVIDUALS WHO HAVE  
FREQUENT CONDOMLESS SEX  
WITH MULTIPLE PARTNERS  
AND ARE AT ELEVATED RISK OF AN STI



Joseph Osmundson, Ian W Holloway, Raul Macias Gil, J Trevor Jann, Chenglin Hong, Robert K Bolan, Doxycycline Postexposure Prophylaxis for Bacterial STIs: Prescribing Patterns, Use, Short-term Outcomes Among 2083 Patients in a Los Angeles Federally Qualified Health Care Program, and Implications for Widespread Use, Open Forum Infectious Diseases, Volume 12, Issue 3, March 2025, ofaf089, <https://doi.org/10.1093/ofid/ofaf089>